

THEREDA REVISITED – PROJECT STATUS IN 2013

V. Brendler⁽¹⁾, F. Bok⁽¹⁾, Ch. Marquardt⁽²⁾, M. Altmaier⁽²⁾

(1) Helmholtz-Zentrum Dresden-Rossendorf, Institute of Resource Ecology, PO Box 510119, D-01314 Dresden, Germany

(2) Karlsruhe Institute of Technology, Institute for Nuclear Waste Disposal, PO Box 3640, D-76021 Karlsruhe, Germany



The Project

- Design of waste repositories requires reliable thermodynamic data to predict the geochemical behaviour of contaminants
- Need: aqueous speciation, solubility limiting solid phases and ion-interaction parameters
- Challenges: Incomplete sets of major and trace elements, inconsistencies between species considered and corresponding formation constants / activity models, restricted variation ranges of parameters (temperature, density, pressure, ionic strength)
- Action: Launch of THEREDA in 2006

The Partners and Goal

- Gesellschaft für Anlagen- und Reaktorsicherheit mbH, Braunschweig (Coordinator)
- Helmholtz-Zentrum Dresden-Rossendorf, Institute of Resource Ecology
- Karlsruhe Institute of Technology, Institute for Nuclear Waste Disposal
- TU Bergakademie Freiberg, Institute of Inorganic Chemistry
- AF-Consult Switzerland AG, Baden (Switzerland).

Main objective: a centrally administrated and maintained database of verified thermodynamic parameters for environmental applications in general and radiochemical issues in particular

The Chemistry

- Actinides, Fission and Activation Products:
 - ◆ Pa, Th, U, Np, Pu, Am, Cm
 - ◆ Rb, Sr, Tc, Cs, Sm, Nd, Ra
- Matrix:
 - ◆ System of oceanic salts: Na⁺, K⁺, H⁺, OH⁻, Mg²⁺, Ca²⁺, Cl⁻, SO₄²⁻, CO₃²⁻ / HCO₃⁻ / CO₂(aq) – H₂O containing the elements Na, K, Mg, Ca, Cl, S, C
 - ◆ Hydrated cement phases (including Al, Si)
- Ion-Ion-Interactions:
 - ◆ Pitzer approach
 - ◆ (SIT & extended Debye-Hückel)

Releases / Extended Interface

- Six new official releases since Migration 2011, all based on Pitzer (in bold: additions to the hexary system of oceanic salts)
 - ◆ Na, Mg, Ca – Cl – **Am(III), Nd(III), Cm(III)** – H₂O(l) (2011-11-23)
 - ◆ Na, K, Mg, Ca – Cl, SO₄ – **HCO₃/CO₂(g)** – H₂O(l) (2012-02-21)
 - ◆ Na – Cl – **Np(V)** – H₂O(l) (2012-11-30)
 - ◆ Na, Mg, Ca, K – Cl, SO₄ – **HCO₃/CO₂(g)** – **Cs** – H₂O(l) (2013-01-28)
 - ◆ Na, K, Mg, Ca – Cl, SO₄ – **HCO₃/CO₂(g)** – **Si, Al** – H₂O(l) (2013-07-04)
 - ◆ Na, K, Ca – Cl – **HCO₃/CO₂(g)** – **Th(IV), Np(IV), Pu(IV)** – H₂O(l) (2013-08-04)
 - ◆ Additional release for U(IV) and U(VI) is planned for the near future.
- Download as separate files (www.thereda.de → navigation menu: THEREDA Data Query → Tailored Databases) in five formats: generic ASCII type and specific to geochemical speciation codes (PhreeqC, EQ3/6, ChemApp, Geochemist's Workbench – GWB).
- Access to data records is now also possible through interactive forms (menu: THEREDA Data Query → Single Data Query // Complex Systems), both with export options as CSV or MS Excel file.
- New interactive web-based tool for data entry and editing – though only visible to members of the editorial board of THEREDA.
- Eight issued technical papers (Downloads → Documentations) add transparency and promote the transition of THEREDA into an open information and discussion platform – covering the database itself as well as geochemical modeling at large.

Quality Assurance (QA)

- Major milestones in QA:
 - ◆ Integration of a new auditing scheme,
 - ◆ Monitor data selections
 - ◆ Automatic logging system storing all data changes in separate files
 - ◆ Extension of the internal calculation scheme for mutually dependent thermodynamic data to check for data consistency and plausibility.
- An illustration of THEREDA's capability: the solubility of freshly precipitated Nd(OH)₃(am) in 3.86 m CaCl₂ solution.
- Calculations compared to predictions (based on two databases from the Yucca Mountain Project – data0.ypf.R1/2) and experimental results (Fig. 1).
- Deviations in the weak alkaline range due to different solubility constants for Nd(III) hydroxides ($\Delta \log K_{\text{Nd(OH)}_3(\text{am})} \approx 3.3$).
- The missing increase of Nd(III) solubility in higher alkaline range ($-\log(\text{mH}^+) > 10$) using the data0.ypf.R1/2 databases is a consequence of different aqueous speciation, especially for anionic and ternary Ca-Nd-(OH)_x complexes.




Fig. 1: Solubility of amorphous/crystalline Nd(OH)₃(am) in 3.86 m CaCl₂ solution (lines: database predictions, symbols: experimental data from Neck, V.; Altmaier, M.; Rabung, T.; Lützenkirchen, J.; Fanghänel, T.; Pure and Applied Chemistry 81, 2009, 1555–1568.).

Surface Complexation

- A holistic view of geochemical processes must include sorption phenomena.
- Thermodynamically consistent treatment: only possible with surface complexation modeling (SCM).
- Support for models implemented in PhreeqC and GWB: Diffuse Double Layer Model (DDL), Constant Capacitance Model (CC), Non-electrostatic Model (NE), 1pK-Basic Stern Model (1pK-BS).
- Stored for each mineral: specific surface area, surface binding site density, surface protolysis constants (pK). Capacitance values will be kept for the CC and the 1pK-BS model. Surface complex formations constants (logK) are entered for given combinations of mineral, ligand, and model.
- Currently SCM data are stored for 298.15 K only, but adaptation to temperature-dependent functions is possible.
- The extension towards SCM required the synchronization of various table designs (e.g. for solids and bibliography) and the introduction of numerous new tables, coupled to new data categories.
- Severe general problems: speculative species, unreliable, contradicting or missing thermodynamic parameters.
- **We are looking forward to receive helpful feedback on THEREDA and are hoping for intensified interactions of the THEREDA database project with the international user community.**

Acknowledgements

This project is jointly funded by the German Federal Ministries of Education and Research (BMBF), of Economics and Technology (BMWi), and for the Environment, Nature Conservation and Nuclear Safety (BMU).